

Armenia

Tuberculosis case notifications, 2006







Total number of cases	2 155
Notification rate per 100 000	71.6
Sex ratio (M:F)	4.0
Median age-group, nationals	35-44 years
Median age-group, non-nationals	-
Foreign citizens	0 (0.0%)
New (never-treated)	1 598 (74.2%)
Culture positive	-
Pulmonary	1 779 (82.6%)
of which sputum smear positive	884 (49.7%)
HIV positive	25 (1.2%)
TB deaths per 100 000 (2003)	4.83

Drug Resistance Surveillance, 2006

Geographic coverage	National
International proficiency testing	Yes (2005)
Case-linked data reporting	No *
Cases with DST results	870
Cases resistant to isoniazid	370 (42.5%)
Cases resistant to rifampicin	234 (26.9%)
MDR cases	215 (24.7%)
Cases resistant to ethambutol	133 (15.3%)
Cases resistant to streptomycin	383 (44.0%)
* Data from NRL (representativeness unknown)	
Culture and DST not routinely performed	

Treatment Outcome Monitoring, 2005

Geographic coverage	National
Outcome cohort	All pulmonary smear positive
Case-linked data reporting	No
Included in TOM cohort	908
Success	554 (61%)
Died	44 (5%)
Failed	68 (7%)
Still on treatment	0 (0%)
Lost to follow up	242 (27%)



† Origin unknown in 2001