

Belarus

Tuberculosis case notifications, 2006

Total number of cases	6 065
Notification rate per 100 000	62.3
Sex ratio (M:F)*	2.5
Median age-group, nationals*	35-44 years
Median age-group, non-nationals	-
Foreign citizens	0 (0.0%)
New (never-treated)	5 142 (84.8%)
Culture positive	2 126 (35.1%)
Respiratory	5 679 (93.6%)
of which sputum smear positive	1 072 (18.9%)
HIV positive new TB cases (2005)	32 (0.6%)
TB deaths per 100 000 (2003)	10.4

Drug Resistance Surveillance, 2000



Geographic coverage	Unknown
International proficiency testing	No
Case-linked data reporting	No †
Cases with DST results	2 060
Cases resistant to isoniazid	-
Cases resistant to rifampicin	-
MDR cases	220 (10.7%)
Cases resistant to ethambutol	-
Cases resistant to streptomycin	-

† Reported as aggregate data on new cases

Treatment Outcome Monitoring, 2005

Geographic coverage	National
Outcome cohort	New pulm smear &/or culture positive
Case-linked data reporting	No
Included in TOM cohort	2 247
Success	1 642 (73%)
Died	228 (10%)
Failed	237 (11%)
Still on treatment	0 (0%)
Lost to follow up	140 (6%)


* For new cases only



§ Retreated cases not included 2003-2005

Tuberculosis cases by geographic origin, 1995-2006

Foreign citizens not reported



* Retreated cases not included

Combined multidrug resistance, by origin, 2001-2006

Not available

