

Poland

Tuberculosis case notifications, 2006

Total number of cases	8 593
Notification rate per 100 000	22.5
Sex ratio (M:F)	2.0
Median age-group (all cases)	45-54 years
Median age-group, non-nationals	25-34 years
Foreign citizens	47 (0.5%)
New (not previously diagnosed)	7 585 (88.3%)
Culture positive	5 233 (60.9%)
Pulmonary	7 884 (91.7%)
of which sputum smear positive	3 310 (42.0%)
HIV positive TB cases (2003)	15 (0.1%)
TB deaths per 100 000 (2005)	2.11







Drug Resistance Surveillance, 2004

Geographic coverage	National
International proficiency testing	Yes (2004)
Case-linked data reporting	No *
Cases with DST results	3 239
Cases resistant to isoniazid	162 (5.0%)
Cases resistant to rifampicin	66 (2.0%)
MDR cases	51 (1.6%)
Cases resistant to ethambutol	16 (0.5%)
Cases resistant to streptomycin	131 (4.0%)

* Survey of all DST laboratories

Treatment Outcome Monitoring, 2005

Geographic coverage	National
Outcome cohort	All pulmonary culture positive
Case-linked data reporting	Yes
Included in TOM cohort	5 178
Success	3 760 (73%)
Died	286 (6%)
Failed	33 (1%)
Still on treatment	21 (0%)
Lost to follow up	1 078 (21%)



† No data by origin except in 2004 survey