

Serbia*

Tuberculosis case notifications, 2006

Total number of cases	2 150 †
Notification rate per 100 000	29.0
Sex ratio (M:F)	1.6
Median age-group, nationals	45-54 years
Median age-group, non-nationals	45-54 years
Foreign citizens	20 (0.9%)
New (not previously diagnosed)	1 868 (86.9%)
Culture positive	1 271 (59.1%)
Pulmonary	1 853 (86.2%)
of which sputum smear positive	984 (53.1%)
HIV positive TB cases	5 (0.2%)
TB deaths per 100 000 (2002)	3.29

Drug Resistance Surveillance, 2006


Geographic coverage	Partial †
International proficiency testing	Yes
Case-linked data reporting	Yes
Cases with DST results	1 269
Cases resistant to isoniazid	21 (1.7%)
Cases resistant to rifampicin	11 (0.9%)
MDR cases	11 (0.9%)
Cases resistant to ethambutol	6 (0.5%)
Cases resistant to streptomycin	31 (2.4%)
† Data representativeness unknown	

Treatment Outcome Monitoring, 2005


Geographic coverage	National	
Outcome cohort	All pulmonary culture positive	
Case-linked data reporting	Yes	
Included in TOM cohort	1 338	
Success	1 120	(84%)
Died	73	(5%)
Failed	15	(1%)
Still on treatment	1	(0%)
Lost to follow up	129	(10%)

† Excluding Kosovo (1,122 cases in 2006)


Tuberculosis notification rates by sex, 1995-2006*




Tuberculosis notification rates by age group, 1995-2006*



Tuberculosis cases by geographic origin, 1995-2006*




Tuberculosis cases by origin, age group and sex, 2006




Combined multidrug resistance, by origin, 2001-2006

Data representativeness unknown



§ No data for 2001; origin unknown in 2002

Outcomes, new pulmonary definite cases, 2001-2005§



§ Belgrade region 2001-2004; nationwide pulmonary culture positive in 2005

* established in 2006 following the split of Serbia & Montenegro; notifications for Kosovo included until 1997 and for Montenegro until 2004